

## International Hospital Inpatient Quality Measures

<b>I-Emergency Department (I-ED) – Stratified (posted April 2018)</b>		
<b>Measure Code</b>	<b>Measure Short Name</b>	<b>Measure Description</b>
<b>I-ED-1a</b>	Median Time from ED Arrival to ED Departure for Admitted ED Patients – Overall Rate	Median time from emergency department arrival to time of departure from the emergency room for patients admitted to the facility from the emergency department – all patients
<b>I-ED-1b</b>	Median Time from ED Arrival to ED Departure for Admitted ED Patients – Reporting Measure	Median time from emergency department arrival to time of departure from the emergency room for patients admitted to the facility from the emergency department – <b>excludes</b> Psychiatric/Mental Health Patients
<b>I-ED-1c</b>	Median Time from ED Arrival to ED Departure for Admitted ED Patients – Psychiatric/Mental Health Patients	Median time from emergency department arrival to time of departure from the emergency room for patients admitted to the facility from the emergency department – <b>includes only</b> Psychiatric/Mental Health patients
<b>I-ED-2a</b>	Admit Decision Time to ED Departure Time for Admitted Patients – Overall Rate	Median time from admit decision time to time of departure from the emergency department for admitted patients. – all patients
<b>I-ED-2b</b>	Admit Decision Time to ED Departure Time for Admitted Patients – Reporting Measure	Median time from admit decision time to time of departure from the emergency department for admitted patients - <b>excludes</b> Psychiatric/Mental Health Patients
<b>I-ED-2c</b>	Admit Decision Time to ED Departure Time for Admitted Patients –Psychiatric/Mental Health Patients	Median time from admit decision time to time of departure from the emergency department for admitted patients - <b>includes only</b> Psychiatric/Mental Health patients

Use **Ctrl + Home** keys to go to the title page

## International Hospital Inpatient Quality Measures

<b>I-Acute Myocardial Infarction (I-AMI)</b>		
<b>Measure Code</b>	<b>Measure Short Name</b>	<b>Measure Description</b>
I-AMI-1	Aspirin at Arrival	Aspirin received within 24 hours of arrival to the hospital for patients having an acute myocardial infarction (AMI).
I-AMI-2	Aspirin Prescribed at Discharge	Aspirin prescribed at discharge for patients who had an acute myocardial infarction.
I-AMI-3	ACEI or ARB for LVSD	ACEI (angiotensin converting enzyme inhibitor) or ARB (angiotensin receptor blocker) for patients who have LVSD (Left Ventricular Systolic Dysfunction) after having an acute myocardial infarction.
I-AMI-4	Adult Smoking Cessation Advice/Counseling	Adult smoking (cigarette) cessation advice/counseling given to patients who had an acute myocardial infarction.
I-AMI-5	Beta Blocker Prescribed at Discharge	Beta-blocker prescribed at discharge for patients who had an acute myocardial infarction.
I-AMI-9	Inpatient Mortality	Acute myocardial infarction (AMI) patients who expire during the hospital stay

<b>I-Heart Failure (I-HF)</b>		
<b>Measure Code</b>	<b>Measure Short Name</b>	<b>Measure Description</b>
I-HF-2	Evaluation of LVS Function	Heart failure patients with documentation in the hospital record that left ventricular systolic (LVS) function was evaluated before arrival; during hospitalization, or is planned for after discharge
I-HF-3	ACEI or ARB for LVSD	ACEI (angiotensin converting enzyme inhibitor) or ARB (angiotensin receptor blocker) for heart failure

Use [Ctrl + Home](#) keys to go to the title page

## International Hospital Inpatient Quality Measures

		patients who have LVSD (Left Ventricular Systolic Dysfunction)
I-HF-4	Adult Smoking Cessation Advice/Counseling	Adult smoking(cigarettes) cessation advice/counseling given to heart failure patients

### I-Pneumonia (I-PN)

Measure Code	Measure Short Name	Measure Description
I-PN-2	Pneumococcal Vaccination	Pneumonia patients, aged 65 and older, who were screened for pneumococcal vaccine status and were administered the vaccine prior to discharge, if indicated
I-PN-4	Adult Smoking Cessation Advice/Counseling	Adult smoking cessation advice/counseling given to patients who smoke cigarettes and who are hospitalized for pneumonia
I-PN-7	Influenza Vaccination	Pneumonia patients, aged 50 and older, who during the flu season, were screened for influenza vaccine status and were vaccinated prior to discharge, if indicated

### I-Surgical Care Improvement Project (I-SCIP)

Measure Code	Measure Short Name	Measure Description
I-SCIP-Inf-1d	Prophylactic Antibiotic Received Within One Hour Prior to Surgical Incision Hip Arthroplasty	<p>Surgical patients with prophylactic antibiotics initiated within one hour prior to surgical incision.</p> <ul style="list-style-type: none"> <li>Patients who received Vancomycin or a Fluroquinolone for prophylactic antibiotics should have the antibiotics initiated within two hours prior to surgical incision.</li> <li>Due to the longer infusion time required for Vancomycin and</li> </ul>

Use **Ctrl + Home** keys to go to the title page

## International Hospital Inpatient Quality Measures

		Fluroquinolone, it is acceptable to start these antibiotics within two hours prior to incision time.
I-SCIP-Inf-1e	Prophylactic Antibiotic Received Within One Hour Prior to Surgical Incision-Knee Arthroplasty	<p>Surgical patients with prophylactic antibiotics initiated within one hour prior to surgical incision.</p> <ul style="list-style-type: none"> <li>Patients who received Vancomycin or a Fluroquinolone for prophylactic antibiotics should have the antibiotics initiated within two hours prior to surgical incision.</li> <li>Due to the longer infusion time required for Vancomycin and Fluroquinolone, it is acceptable to start these antibiotics within two hours prior to incision time.</li> </ul>
I-SCIP-Inf-2d	Prophylactic Antibiotic Selection for Surgical Patients-Hip Arthroplasty	<p>Surgical patients who received prophylactic antibiotics consistent with current Hip Arthroplasty guidelines,</p> <ul style="list-style-type: none"> <li>Appendix C, Table 3.2, Prophylactic Antibiotic Regimen Selection for Surgery</li> </ul>
I-SCIP-Inf-2e	Prophylactic Antibiotic Selection for Surgical Patients-Knee Arthroplasty	<p>Surgical patients who received prophylactic antibiotics consistent with current Knee Arthroplasty guidelines</p> <ul style="list-style-type: none"> <li>Appendix C, Table 3.2, Prophylactic Antibiotic Regimen Selection for Surgery</li> </ul>
I-SCIP-Inf-3d	Prophylactic Antibiotics Discontinued Within 24 Hours After Surgery End Time- Hip Arthroplasty	Surgical patients, who had a Hip Arthroplasty, whose prophylactic antibiotics were discontinued within 24 hours after <i>Anesthesia End Time</i>

Use [Ctrl + Home](#) keys to go to the title page

## International Hospital Inpatient Quality Measures

I-SCIP-Inf-3e	Prophylactic Antibiotics Discontinued Within 24 Hours After Surgery End Time Knee Arthroplasty	Surgical patients, who had a Knee Arthroplasty, whose prophylactic antibiotics were discontinued within 24 hours after <i>Anesthesia End Time</i>
I-SCIP-VTE-1	Surgery Patients with Recommended Venous Thromboembolism Prophylaxis <u>Ordered</u>	Surgery patients with recommended Venous Thromboembolism (VTE) prophylaxis <u>ordered</u> anytime from <i>hospital arrival</i> to 24 hours after <i>Anesthesia End Time</i> ,
I-SCIP-VTE-2	Surgery Patients Who <u>Received</u> Appropriate Venous Thromboembolism Prophylaxis Within 24 hours Prior to Surgery to 24 Hours After Surgery	Surgery patients who received appropriate Venous Thromboembolism (VTE) prophylaxis within 24 hours prior to <i>Anesthesia Start Time</i> to 24 hours after <i>Anesthesia End Time</i> ,

### I-Hospital Based Inpatient Psychiatric Care Services (I-HBIPS)

Measure Code	Measure Short Name	Measure Description
I-HBIPS-2	Hours of physical restraint use	The total number of hours that all patients admitted to a hospital-based inpatient psychiatric setting were maintained in physical restraint
I-HBIPS-3	Hours of seclusion use	The total number of hours that all patients admitted to a hospital-based inpatient psychiatric setting were held in seclusion.

### I-Children's Asthma Care (I-CAC)

Measure Code	Measure Short Name	Measure Description
I-CAC-1	Relievers for Inpatient Asthma	Use of relievers in pediatric patients admitted for inpatient treatment of asthma
I-CAC-2	Systemic Corticosteroids for Inpatient Asthma	Use of systemic corticosteroids in pediatric patients admitted for inpatient treatment of asthma

Use [Ctrl + Home](#) keys to go to the title page

## International Hospital Inpatient Quality Measures

<b>I-Venous Thromboembolism (I-VTE)</b>		
<b>Measure Code</b>	<b>Measure Short Name</b>	<b>Measure Description</b>
I-VTE-1	Venous Thromboembolism Prophylaxis	This measure assesses the number of patients who received VTE prophylaxis or have documentation why no VTE prophylaxis was given the day of or the day after hospital admission or surgery end date for surgeries that start the day of or the day after hospital admission.
I-VTE-2	Intensive Care Unit Venous Thromboembolism Prophylaxis	This measure assesses the number of patients who received VTE prophylaxis or have documentation why no VTE prophylaxis was given the day of or the day after the initial admission (or transfer) to the Intensive Care Unit (ICU) or surgery end date for surgeries that start the day of or the day after ICU admission (or transfer).

<b>I-Stroke (I-STK)</b>		
<b>Measure Code</b>	<b>Measure Short Name</b>	<b>Measure Description</b>
I-STK-2	Discharged on Antithrombotic Therapy	Ischemic stroke patients prescribed antithrombotic therapy at hospital discharge.
I-STK-3	Anticoagulation Therapy for Atrial Fibrillation/Flutter	Ischemic stroke patients with atrial fibrillation/flutter who are prescribed anticoagulation therapy at hospital discharge.
I-STK-8	Stroke Education	Ischemic or hemorrhagic stroke patients or their caregivers who were given educational material addressing ALL of the following: Activation of emergency medical system (if available in region), need for follow-up after discharge, medications prescribed at discharge,

Use [Ctrl + Home](#) keys to go to the title page

## International Hospital Inpatient Quality Measures

		risk factors for stroke, and warning signs and symptoms of stroke.
I-STK-10	Assessed for Rehabilitation	Ischemic or hemorrhagic stroke patients who were assessed for or received rehabilitation services.

<b>I-Nursing- Sensitive Care (I-NSC)</b>		
<b>Measure Code</b>	<b>Measure Short Name</b>	<b>Measure Description</b>
I-NSC-2	Pressure Ulcer Prevalence (Hospital-Acquired)	Patients that have hospital-acquired (nosocomial) category/stage II or greater pressure ulcer(s) on the day of the prevalence study.
I-NSC-4	Patient Falls	All documented falls with or without injury, experienced by patients in a calendar month.
I-NSC-5	Falls with Injury	All documented falls by a patient with an injury level of “minor” (2) or greater.

<b>I-Perinatal Care (I-PC)</b>		
<b>Measure Code</b>	<b>Measure Short Name</b>	<b>Measure Description</b>
I-PC-1	Elective Delivery	Patients with elective vaginal deliveries or elective cesarean sections at $\geq 37$ and $< 39$ weeks of gestation completed
I-PC-2	Cesarean Section	Nulliparous women with a term, singleton baby in a vertex position delivered by cesarean section
I-PC-5	Exclusive Breast Feeding	Exclusive breast milk feeding during the newborn's entire hospitalization

Use [Ctrl + Home](#) keys to go to the title page