

May 29, 2020

## **Position Statement: Preventing Nosocomial COVID-19 Infections as Organizations Resume Regular Care Delivery**

The Joint Commission supports the following positions for healthcare organizations to prevent nosocomial COVID-19 infections as they are resuming routine care.

**Healthcare organizations should continue to follow [CDC recommendations](#) for universal masking of staff, patients, and visitors.** If there are situations where a patient cannot wear a mask (e.g., under 2 years of age, respiratory compromise, or examination of the nose, mouth, lips, and perioral area) personnel providing care within 6 feet of the patient should don a medical mask. In areas where there is moderate to substantial transmission of COVID-19 in the community (or as indicated by [standard precautions](#)), personnel should also wear eye protection in addition to wearing a mask. If there are no COVID-19 cases in the community for several weeks, organizations should work with public health authorities to re-evaluate the need for universal masking based on their community's risk of new cases; if the organization stops universal masking, it should be prepared to immediately re-institute universal masking if new cases emerge.

**Healthcare personnel should wear filtering facepiece respirators (e.g., N95 respirators) or higher-level respirator for all [aerosol generating procedures](#) or surgical procedures that might pose higher risk for transmission** (e.g., that generate potentially infectious aerosols or involving anatomic regions where viral loads might be higher, such as the nose and throat, oropharynx, respiratory tract). This should be done even if the patient has had a negative test for active COVID-19 infection because of the risk of false-negatives.

**Healthcare organizations should no longer be operating under crisis standards of care and no longer following [contingency strategies for use of gowns, eye protection, or facemasks](#) when they resume elective procedures and ambulatory care.**

In areas that experienced a large surge in the number of COVID-19 cases, many hospitals and healthcare organizations were unable to follow their usual policies and procedures and had to resort to "[crisis standards of care](#)." The effectiveness of crisis strategies is uncertain, and they may pose a risk for transmission of infectious diseases between healthcare providers and patients or other safety concerns. Therefore, the volume of care delivered under crisis standards should be limited, and an expansion of services to elective procedures and ambulatory care would be inappropriate until patient care activities are back within routine standards.

**During times of respirator shortages, it is acceptable for healthcare organizations to use contingency conservation strategies for filtering facepiece respirators (e.g., N95 respirators) to ensure an adequate supply of respirators for all [aerosol-generating procedures](#).**

Re-opening for elective procedures and following recommendations highlighted in this document will greatly increase the demand for respirators. To ensure adequate supplies, organizations may need to continue to use approved [conservation strategies](#). Failure to use conservation strategies could limit the availability of respirators for those situations in which their use is critical to protect healthcare workers.

**Patients undergoing elective surgery (or other [aerosol-generating procedures](#)) should be tested for COVID-19 shortly before the scheduled procedure.**

We support the recommendations of the [American Society of Anesthesiologist and the Anesthesia Patient Safety Foundation](#) for preoperative screening and [testing for current COVID-19 infection](#). We recognize that some organizations will not have adequate testing supplies to accomplish this currently. This is a serious concern; ensuring adequate supplies to test all patients before elective surgery should be a national priority to protect patients and healthcare workers.

Prior to testing, patients should be screened for COVID-19 symptoms; patients with COVID-19 symptoms should be sent immediately for evaluation. When possible, there should be universal testing for COVID-19 among patients without symptoms. This is necessary because patients with incubating COVID-19 could be asymptomatic immediately prior to an elective procedure and then develop overt COVID-19 infection. If the infection is severe, this could lead to [complications during the post-procedure period](#). Moreover, if patients with asymptomatic or presymptomatic COVID-19 undergo surgery, this poses a high risk of transmission of COVID-19 to the operative team. Patients who are found to have active disease (positive COVID-19 test) should not undergo their planned procedure until the infection has resolved.

Ideally, testing should be done as close as possible to the time of the procedure while ensuring there is adequate time to obtain the results. Some states require testing to be done within 72-96 hours of the procedure. Patients should be advised to self-quarantine between the time of testing and the day of their procedure to minimize the chance of becoming infected during the window between testing and the procedure.

**Patients undergoing elective surgery (or other aerosol-generating procedures) whose preoperative test for COVID-19 is negative should still be treated with universal respiratory precautions because of the risk of false-negative COVID-19 tests.**

Because of sampling technique, sample site, the test performed and the timing of the test, there is a risk of false negative COVID-19 tests. Therefore, we support the recommendations of the [American Society of Anesthesiologist and the Anesthesia Patient Safety Foundation](#) and [others](#) who call for use of a surgical mask and eye covering by OR staff for all operative cases and an N95 mask, eye protection, gloves and a gown to be worn by all in the operating room staff during aerosol-generating procedures (AGP) such as intubation and extubation, surgical procedures that might pose higher risk for transmission such as laparoscopic surgery and gastrointestinal endoscopy that requires insufflation, or procedures in anatomic regions where viral loads might be higher, such as the nose and throat, oropharynx, respiratory tract. If it is not possible to have all staff who are in the room during and after an AGP wear an N95 mask, then staff not wearing an N95 mask may enter the room following an AGP after the [CDC-recommended time for air clearance](#) (99% efficiency) to occur within the room.

**Organizations should check with state and local health departments for more specific requirements.**

For example, some states have established requirements regarding the type of pre-operative/pre-procedure testing for SARS-CoV-2 that must be done and the time period in which it must be done (e.g., no more than three days prior to the surgery or procedure).